



# DeSoto County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Middle School				Comparison of Two Rates		
	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>4.9</b>	(2.6-7.1)	<b>2.3</b>	(2.1-2.5)	+		
Smoked cigars on one or more of the past 30 days	<b>2.7</b>	(0.3-5.2)	<b>2.9</b>	(2.6-3.2)			+
Used smokeless tobacco on one or more of the past 30 days	<b>4.0</b>	(1.8-6.1)	<b>2.1</b>	(1.9-2.3)			+
Used electronic cigarettes on one or more of the past 30 days	<b>3.4</b>	(1.6-5.2)	<b>4.0</b>	(3.6-4.3)			+
Used hookah on one or more of the past 30 days	<b>1.5</b>	(0.0-2.9)	<b>3.5</b>	(3.1-3.9)		+	
Used any form of tobacco on one or more of the past 30 days*	<b>9.2</b>	(5.7-12.7)	<b>5.1</b>	(4.7-5.4)	+		
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	<b>8.9</b>	(5.4-12.5)	<b>8.1</b>	(7.6-8.6)			+
Using electronic cigarettes is less harmful than smoking cigarettes	<b>64.9</b>	(55.2-74.6)	<b>64.7</b>	(63.2-66.1)			+
Using hookah is less harmful than smoking cigarettes	<b>14.3</b>	(6.6-21.9)	<b>42.0</b>	(39.9-44.0)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>68.4</b>	(63.5-73.4)	<b>72.6</b>	(71.7-73.5)			+
Exposed to secondhand smoke during the past 7 days (in a room or car)	<b>39.1</b>	(33.2-45.0)	<b>35.9</b>	(35.0-36.7)			+
Smoking is allowed in the home	<b>11.5</b>	(7.5-15.4)	<b>8.8</b>	(8.3-9.3)			+

\*This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.



# DeSoto County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Middle School				Comparison of Two Rates		
	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Students who were taught about tobacco use in any classes this year	<b>51.1</b>	(44.2-58.0)	<b>47.1</b>	(45.7-48.5)			+
Parent or guardian has talked about the dangers of tobacco in the past year	<b>66.6</b>	(61.0-72.2)	<b>67.5</b>	(66.7-68.3)			+
Served as an active member of SWAT	<b>4.1</b>	(2.2-5.9)	<b>2.9</b>	(2.7- 3.2)			+
Definitely did not think that smokers have more friends	<b>27.5</b>	(21.7-33.4)	<b>31.7</b>	(30.8-32.5)			+
Definitely did not think that smoking helps young people "look cool" or "fit in"	<b>67.6</b>	(62.6-72.7)	<b>70.2</b>	(69.4-70.9)			+
Lifetime asthma	<b>17.6</b>	(13.7-21.5)	<b>20.2</b>	(19.5-20.9)			+
Current asthma	<b>10.8</b>	(6.7-14.8)	<b>12.1</b>	(11.5-12.6)			+
Asthma attack in past year	<b>16.9</b>	(0.6-33.3)	<b>20.0</b>	(18.5-21.5)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>14.6</b>	(9.5-19.7)	<b>12.4</b>	(11.7-13.2)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	<b>20.3</b>	(16.3-24.2)	<b>16.3</b>	(15.6-17.0)			+
Physically active for at least 60 minutes per day during the past 7 days	<b>26.8</b>	(19.7-33.9)	<b>24.7</b>	(24.0-25.6)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>53.0</b>	(47.8-58.2)	<b>44.6</b>	(43.8-45.5)	+		
Described themselves as slightly or very overweight	<b>38.0</b>	(31.8-44.2)	<b>30.1</b>	(29.3-30.8)	+		



# DeSoto County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	High School				Comparison of Two Rates		
	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>12.3</b>	(9.4-15.2)	<b>7.5</b>	(7.1-7.9)	+		
Smoked cigars on one or more of the past 30 days	<b>13.7</b>	(10.0-17.4)	<b>9.1</b>	(8.6-9.6)	+		
Used smokeless tobacco on one or more of the past 30 days	<b>12.7</b>	(9.3-16.2)	<b>5.4</b>	(5.0-5.8)	+		
Used electronic cigarettes on one or more of the past 30 days	<b>12.5</b>	(8.6-16.4)	<b>10.8</b>	(10.3-11.4)			+
Used hookah on one or more of the past 30 days	<b>5.2</b>	(2.7-7.8)	<b>11.6</b>	(11.0-12.1)		+	
Used any form of tobacco on one or more of the past 30 days*	<b>27.6</b>	(23.2-32.1)	<b>14.8</b>	(14.2-15.5)	+		
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	<b>30.0</b>	(25.0-35.0)	<b>22.6</b>	(21.8-23.3)	+		
Using electronic cigarettes is less harmful than smoking cigarettes	<b>69.8</b>	(61.8-77.8)	<b>71.1</b>	(69.9-72.3)			+
Using hookah is less harmful than smoking cigarettes	<b>42.6</b>	(34.5-50.7)	<b>57.3</b>	(56.0-58.6)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>49.4</b>	(44.4-54.3)	<b>60.7</b>	(59.8-61.6)		+	
Exposed to secondhand smoke during the past 7 days (in a room or car)	<b>46.6</b>	(41.1-52.1)	<b>39.7</b>	(38.8-40.6)	+		
Smoking is allowed in the home	<b>13.4</b>	(9.4-17.4)	<b>8.8</b>	(8.3-9.3)	+		

\*This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.



# DeSoto County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	High School				Comparison of Two Rates		
	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Students who were taught about tobacco use in any classes this year	<b>40.9</b>	(32.9-48.9)	<b>29.6</b>	(28.4-30.7)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	<b>49.3</b>	(44.3-54.4)	<b>56.7</b>	(55.8-57.5)		+	
Served as an active member of SWAT	<b>3.7</b>	(1.3-6.2)	<b>3.1</b>	( 2.8- 3.4)			+
Definitely did not think that smokers have more friends	<b>25.7</b>	(21.0-30.3)	<b>25.3</b>	(24.6-26.1)			+
Definitely did not think that smoking helps young people “look cool” or “fit in”	<b>67.9</b>	(62.0-73.7)	<b>67.1</b>	(66.3-67.9)			+
Lifetime asthma	<b>20.1</b>	(16.1-24.1)	<b>21.2</b>	(20.5-21.9)			+
Current asthma	<b>10.7</b>	(7.7-13.6)	<b>10.8</b>	(10.2-11.3)			+
Asthma attack in past year	<b>20.2</b>	(9.4-30.9)	<b>16.5</b>	(15.2-17.7)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>13.9</b>	(9.6-18.3)	<b>12.3</b>	(11.7-12.9)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	<b>21.6</b>	(17.3-25.9)	<b>15.5</b>	(14.9-16.1)	+		
Physically active for at least 60 minutes per day during the past 7 days	<b>25.2</b>	(20.3-30.1)	<b>21.5</b>	(20.8-22.2)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>49.9</b>	(44.4-55.4)	<b>42.5</b>	(41.7-43.4)	+		
Described themselves as slightly or very overweight	<b>39.6</b>	(34.2-45.0)	<b>31.2</b>	(30.4-31.9)	+		



# DeSoto County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>7.3</b>	(5.4-9.1)	<b>4.3</b>	(4.1-4.6)	+		
Smoked cigars on one or more of the past 30 days	<b>7.1</b>	(4.7-9.5)	<b>5.4</b>	(5.2-5.7)			+
Used smokeless tobacco on one or more of the past 30 days	<b>7.2</b>	(4.9-9.4)	<b>3.3</b>	(3.1-3.5)	+		
Used electronic cigarettes on one or more of the past 30 days	<b>7.5</b>	(5.0-9.9)	<b>7.2</b>	(6.8-7.5)			+
Used hookah on one or more of the past 30 days	<b>2.5</b>	(1.1-4.0)	<b>7.1</b>	(6.7-7.5)		+	
Used any form of tobacco on one or more of the past 30 days*	<b>16.2</b>	(12.9-19.6)	<b>9.2</b>	(8.8-9.6)	+		
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	<b>17.0</b>	(13.2-20.7)	<b>14.8</b>	(14.2-15.3)			+
Using electronic cigarettes is less harmful than smoking cigarettes	<b>69.3</b>	(63.1-75.4)	<b>69.4</b>	(68.4-70.3)			+
Using hookah is less harmful than smoking cigarettes	<b>32.5</b>	(25.2-39.8)	<b>52.8</b>	(51.7-54.0)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>60.8</b>	(56.8-64.9)	<b>67.1</b>	(66.4-67.7)		+	
Exposed to secondhand smoke during the past 7 days (in a room or car)	<b>43.6</b>	(39.2-48.0)	<b>37.5</b>	(36.9-38.2)	+		
Smoking is allowed in the home	<b>12.5</b>	(9.4-15.5)	<b>8.4</b>	(8.0-8.7)	+		

\*This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.



# DeSoto County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Students who were taught about tobacco use in any classes this year	<b>46.6</b>	(41.2-52.1)	<b>38.0</b>	(37.1-39.0)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	<b>59.3</b>	(54.9-63.7)	<b>62.4</b>	(61.8-63.0)			+
Served as an active member of SWAT	<b>3.6</b>	(2.0-5.2)	<b>2.8</b>	(2.7- 3.0)			+
Definitely did not think that smokers have more friends	<b>26.3</b>	(22.5-30.2)	<b>28.2</b>	(27.6-28.8)			+
Definitely did not think that smoking helps young people “look cool” or “fit in”	<b>68.6</b>	(64.9-72.3)	<b>68.8</b>	(68.2-69.3)			+
Lifetime asthma	<b>18.1</b>	(15.1-21.2)	<b>20.8</b>	(20.3-21.3)			+
Current asthma	<b>10.4</b>	(7.7-13.1)	<b>11.5</b>	(11.1-11.9)			+
Asthma attack in past year	<b>20.5</b>	(10.2-30.9)	<b>17.7</b>	(16.7-18.7)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>14.2</b>	(10.8-17.5)	<b>12.2</b>	(11.8-12.7)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	<b>21.3</b>	(18.2-24.4)	<b>15.9</b>	(15.4-16.4)	+		
Physically active for at least 60 minutes per day during the past 7 days	<b>24.5</b>	(20.2-28.7)	<b>22.7</b>	(22.2-23.3)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>52.7</b>	(48.8-56.7)	<b>43.7</b>	(43.1-44.4)	+		
Described themselves as slightly or very overweight	<b>40.6</b>	(36.2-45.1)	<b>30.7</b>	(30.2-31.3)	+		